

Pre-Treatment Preparations

- Arrive early and allow 45-60 minutes for numbing of topical anesthetic.
- Avoid skin irritation or intentional skin tanning. Sun-screen is advisable when outdoors during daylight hours.
- Discontinue any irritant topical agents for 2-3 days prior to treatment.
- Arrive for treatment with clean skin. There should be no lotion, make-up, perfume, powder or bath/shower oil present on the skin in the area to be treated.

Avoid Treatment If You Have Any of the Following:

- Pacemaker or internal defibrillator, or other metallic or electronic implant anywhere in the body. The Hand piece should be used at least 1cm away from cochlear implants in the ear.
- Permanent implant in the treated area such as metal plates, screws and metal piercing or silicon, unless deep enough in the periosteal plane.
- Intra-dermal or superficial sub-dermal areas injected with Botox®/HA/collagen/fat injections or other augmentation methods with bio-material, before the product has been dissipated (up to 6 months), except Botox after binding to the facial muscles (3-7 days). It is possible to treat sooner over injectable products placed in the deep, periosteal plane, as soon as the area has healed (1-3 weeks).
- Current or history of skin cancer, or any other type of cancer, or pre-malignant moles.
- Pregnancy and nursing.
- Severe concurrent conditions, such as cardiac disorders or sensory disturbances.
- Impaired immune system due to immunosuppressive diseases such as AIDS and HIV, or use of immunosuppressive medications.
- Patients with history of diseases stimulated by heat, such as recurrent Herpes Simplex in the treatment area, may be treated only following a prophylactic regime.
- Poorly controlled endocrine disorders, such as diabetes or thyroid dysfunction and hormonal virilization.
- Any active skin condition in the treatment area, such as sores, psoriasis, eczema, and rash.
- History of skin disorders, keloids, abnormal wound healing, as well as very dry and fragile skin.

- History of bleeding coagulopathies or use of anticoagulants in the last 10 days
- Any facial surgery performed within a year prior to treatment.
- Facial dermabrasion, facial resurfacing, or deep chemical peeling within the last three months, if face is treated.
- Having received treatment with light, laser, RF, or other devices in the treated area within 2-3 weeks for non-ablative procedures, and 6-12 weeks for ablative fractional laser resurfacing (according to treatment severity) prior to treatment, except special recommendations.
- Use of Isotretinoin (Accutane®) within 6 months prior to treatment.
- Use of non-steroidal anti-inflammatory drugs (NSAIDS, e.g., ibuprofen-containing agents) one week before and after each treatment session, as per the practitioner's discretion.
- Treating over tattoo or permanent makeup to be kept.
- Treating over the lips.
- Skin type VI and dark VI patients treat with caution.
- Treating over hair bearing surfaces.
- Irritable skin like excessively tanned skin from sun, tanning beds or tanning creams and sprays within the last two weeks.
- As per the practitioner's discretion, we may refrain from treating you if you have any condition that might make it unsafe.